



**Faye Barthold** 

May 4, 2010





### Introduction

### Purpose

- Accelerate the assessment and implementation of new technology, research results, and other scientific advancements from the research and development communities to enhance HPC products and services
- Funded through NCEP and NOAA HMT
- Staffing
  - One full time contractor
  - One FTE 90%
  - Support from HPC DTB staff
- Location
  - Collocated with HPC operations







## **2009 Accomplishments**

- Introduced rime factor tool to HPC forecasters
- Completed a comparison of the 1993 and 2008 Midwest floods
- Investigated predecessor rainfall events (PREs)






- What is rime factor?
  - Indicates amount of riming present on an individual ice particle
  - Values range from 1 (unrimed snow) to >40 (sleet)
  - Instantaneous output
- Ice accumulation rate
  - Derived from the rime factor based on an assumption of a maximum possible snow to liquid ratio
- Available guidance
  - Hourly rime factor and ice accumulation rate forecasts from 00Z and 12Z NAM
  - Estimated 6hr, 12hr, and 24hr total accumulations






# **NAM Rime Factor Example**







# **NAM Rime Factor Example**








- Rainfall during the 13 day periods of 29 June-11 July 1993 and 1-14 June 2008 resulted in severe flooding over the central US
  - Represent the two wettest 13 day periods on record in the warm season based on a 60 year climatology
  - Both events preceded by a wet late winter/spring, resulting in high antecedent soil moisture
- Events characterized by similar patterns in 500 hPa geopotential heights, 250 hPa winds, 850 hPa v-winds, and mean sea level pressure
  - 0.939 correlation between 500 hPa height anomalies between events






### Precipitation and 500mb Height Anomalies







#### Moisture Flux



90 110 130 150 170 190 210 230 250 270 300 330 360

90 110 130 150 170 190 210 230 250 270 300 330 360








# Predecessor Rainfall Events (PREs)

- Coherent area of rain located ahead of a tropical cyclone
  - Rainfall rates > 100mm in 24hr
  - Moisture transport between tropical cyclone and PRE
- Represent an enhanced flooding risk
  - Bring heavy rain to areas outside the path of the tropical cyclone
  - Produce heavy precipitation in advance of the precipitation associated with the tropical cyclone










## **Predecessor Rainfall Events**

## Mo PR Treance











- Object oriented verification (MODE)
  - May provide information about model and forecaster biases in specific precipitation regimes
  - Output will be a component of the on-site HPC QPF experiment
- HWT Spring Experiment—QPF component
  - Goal—evaluate the use of high resolution model and ensemble data for QPF forecasting and identify the challenges associated with using this data





## **Questions?**

Faye.Barthold@noaa.gov Mike.Bodner@noaa.gov Edwin.Danaher@noaa.gov






# **Backup Slides**





### Precipitation and 500mb Heights—1993







### Precipitation and 500mb Heights—2008

